

The intriguing case of Sk-69°194

Pablo G. Ostrov*

Facultad de Ciencias Astronómicas y Geofísicas, Paseo del Bosque S/N (1900) La Plata, Argentina

ABSTRACT

In this paper we report the discovery of eclipses in the early-type Magellanic binary Sk-69°194. We derive an ephemeris for this system, and present a CCD V light curve together with CCD spectroscopic observations. We also briefly discuss the nature of the binary components.

Key words: binaries: eclipsing – stars: early-type – stars: fundamental parameters – stars: individual: Sk-69°194

1 INTRODUCTION

Sk-69°194 ($\alpha = 5 : 34 : 36$, $\delta = -69 : 45 : 37$, J2000) is one of the brightest stars in the OB association LH 81 (Lucke & Hodge 1970), in the Large Magellanic Cloud. Westerlund (1961) studied that region, and obtained for Sk-69°194 (which he identified as the star number 10 in Table 28) $V = 12.30$ and $B - V = -0.15$ (notice that Figs. 7 and 8 are interchanged in Westerlund's work, i.e. Sk-69°194 is the star number 10 in his Fig. 7). In its regard, he just pointed out that probably it was a variable star. The name Sk-69°194 comes from the Sanduleak's objective prism survey (Sanduleak 1969). Later on, Isserstedt (1975) performed UBV photo-electric photometry, obtaining $V = 11.98$; $B - V = -0.08$ and $U - B = -0.96$. He also suspected that Sk-69°194 was a photometric variable. More recently, Massey et al. (2000) classified Sk-69°194 as B0I+WN.

LH 81 is one of the regions that we observed when searching new eclipsing binaries. In this paper we announce that Sk-69°194 is an eclipsing binary system and present its light curve and radial velocity measurements. We have also tried to model our data by means of the Wilson-Devinney (hereafter W-D) programs (Wilson & Devinney 1971, Wilson 1990, Wilson 1993). Based on the spectroscopic data and the results of the W-D model, we discuss the nature of the binary components.

2 OBSERVATIONS AND REDUCTIONS

CCD V photometry of Sk-69°194 was acquired during four observing runs between 1998 and 2001, with the 2.15-m telescope at CASLEO¹. A focal reducer was used, yielding a circular field of diameter about 9 arcmin and a scale of 0.813

arcsec px^{-1} . The reduction of the data was performed by means of IRAF² routines.

The CCD spectrograms were obtained with a REOSC³ spectrograph used in its single dispersion mode, attached to the same telescope between 2000 and 2001. A 600 grooves mm^{-1} grating was used, giving a reciprocal dispersion of $1.63 \text{ \AA px}^{-1}$. Our spectrograms comprise the range from about 3900 to 5500 $\text{\AA}$. The reductions and analysis of the spectroscopic observations were also performed by means of IRAF.

Both the photometric and the spectroscopic observations were acquired by means of a CCD Tek 1024 $\times$ 1024 detector.

3 PHOTOMETRY

Aperture photometry was carried out using a stand-alone version of DAOPHOT (Stetson 1987, 1991). In order to tie all the instrumental magnitudes to a common instrumental system, we used about 15 stars in the field of Sk-69°194 as local standards. Just after the first observing run, the photometric data showed deep eclipses, of the order of a half magnitude.

During a photometric night, in December 5, 1998, we also acquired B and R frames of the Sk-69°194 field, together with photometric standards of the selected areas SA-92 and SA-98 (Landolt 1992), with the aim of linking the instrumental and standard photometric systems.

Técnicas de la República Argentina and the National Universities of La Plata, Córdoba and San Juan.

² IRAF software is distributed by NOAO, operated by AURA for NSF.

³ The REOSC spectrograph is on long term loan from the Liège Observatory.

* E-mail: ostrov@fcaglp.edu.ar

¹ Complejo Astronómico El Leoncito, operated under agreement between the Consejo Nacional de Investigaciones Científicas y




Figure 1. The observed and modelled V light curves for Sk-69°194 and their concomitant O-C residuals. Circles, triangles, squares and diamonds stand for the data acquired during the observing runs in 1998, 1999, 2000 and 2001, respectively.

From the observations of that night, we derive the following magnitudes and colours for Sk-69°194 :

$$V = 12.378 \pm 0.010$$

$$B - V = 0.013 \pm 0.016$$

$$V - R = -0.010 \pm 0.014$$

at HJD=2451153.813. This corresponds to $\phi = 0.534$, therefore the colour indexes must be considered with caution. Besides, Sk-69°194 showed to have slight variations in the shape of its eclipses (see below). The magnitude corresponding to $\phi = 0.25$ is $V = 11.83$. Table 1 displays our V aperture photometry (transformed to the standard system), together with the internal errors derived from the local standards, seeing and airmass.

4 SPECTROSCOPY

The spectrum of Sk-69°194 exhibits noticeable changes with phase. Only the spectrum of one of the stars is distinguishable. Based on our data and ultraviolet observations downloaded from the IUE⁴ archive, we classified the spectrum according to the classification criteria of Walborn & Fitzpatrick (1990), corresponding approximately to the type BN0 Ia. That spectrum is easily measurable, and shows orbital velocity excursions of $\sim 425 \text{ km s}^{-1}$ of amplitude. The

Table 2. Radial velocities

HJD-2450000	phase [km s ⁻¹]	B0I [km s ⁻¹]
2253.711	0.504	170 ± 21
2286.613	0.195	482 ± 30
2287.704	0.284	429 ± 23
2293.633	0.769	82 ± 26
2296.655	0.016	246 ± 16
2328.651	0.634	106 ± 20

emission in the region around NIII 4634–HeII 4686, very weak compared with which is usual in WN stars (as also pointed out Massey et al. 2000), does not show any measurable velocity variation as far as can be determined from our data. This spectral region also shows the more striking phase changes, due to the blend of the emission with the supergiant’s absorption lines.

We determined the radial velocity at phase 0.016 (when the B supergiant is at inferior conjunction) measuring several absorption lines. The velocities corresponding to the other phases were derived by cross correlation by means of the FXCOR IRAF task, using the spectra taken at phase 0.016 as template and avoiding the range of 4500–4800 Å. The measured velocities are given in Table 4.

Four of the obtained spectra, corresponding to both conjunctions and both quadratures, are displayed in Fig. 2. It is interesting to point out that the NIII 4515 absorption is more notorious in the spectra acquired near quadratures.

5 EPHEMERIS

It was not straightforward to derive an ephemeris only from the photometric observations. This was due to the fact that the shape of the light curve varied between 1998 and 2001, and to the intrinsic incompleteness of our data. With the aid of our radial velocity measurements, we succeeded in deriving a period and a time of minimum,

$$P = 12.2252 \pm 0.0035$$

$$T_0 = 2451501.815 \pm 0.012$$

The phase-folded light curve is displayed in Fig. 1. The data obtained during different observing seasons are plotted using different symbols, in order to show the shape variation. We emphasize that the data corresponding to secondary eclipse (observed during 1998 as well as during 2000) was obtained, in both cases, during consecutive nights; i.e. the discrepancy can not be ascribed to a period uncertainty.

6 LIGHT AND VELOCITY CURVE MODELLING

Being visible the spectrum of only one binary component, the derivation of masses and absolute dimensions for this system is not straightforward. However, an alternative approach suitable when the system present a Roche-lobe filling geometry and deep eclipses is to obtain the mass ratio solely from the light curve. We attempted to follow that approach,

⁴ Operated by the National Aeronautics and Space Administration, the European Space Agency and the Science and Engineering Research Council of the U.K.

Table 1. V light photometry of Sk-69°194

HJD 2450000+	V	σ_1	fwhm "	X	HJD 2450000+	V	σ_1	fwhm "	X
1149.859	11.881	0.011	1.98	1.44	1504.621	11.883	0.007	2.76	1.45
1151.605	11.944	0.007	1.96	1.41	1504.729	11.873	0.007	3.27	1.27
1151.663	11.973	0.008	1.92	1.30	1504.768	11.868	0.007	2.68	1.27
1151.706	11.968	0.008	1.94	1.27	1504.806	11.864	0.005	3.12	1.29
1151.773	11.988	0.006	2.16	1.29	1504.834	11.861	0.007	2.96	1.33
1151.844	12.005	0.005	2.09	1.41	1859.590	11.849	0.007	3.42	1.68
1152.655	12.235	0.003	2.13	1.31	1859.607	11.843	0.010	3.50	1.60
1152.701	12.249	0.009	1.83	1.27	1859.650	11.848	0.007	2.78	1.45
1152.744	12.272	0.007	2.13	1.27	1859.685	11.849	0.009	3.14	1.36
1152.793	12.288	0.008	2.19	1.32	1859.707	11.855	0.007	3.70	1.32
1152.824	12.304	0.005	2.22	1.37	1859.740	11.852	0.028	5.70	1.29
1152.860	12.313	0.009	2.06	1.46	1859.743	11.848	0.011	4.68	1.28
1153.678	12.429	0.005	1.56	1.28	1859.778	11.852	0.008	5.36	1.27
1153.729	12.411	0.005	1.69	1.27	1859.803	11.842	0.008	4.80	1.27
1153.761	12.398	0.006	1.77	1.29	1859.841	11.838	0.016	4.53	1.30
1153.811	12.378	0.006	2.40	1.35	1859.867	11.847	0.008	4.51	1.33
1153.852	12.366	0.004	2.64	1.45	1860.610	11.907	0.009	4.99	1.58
1154.699	12.111	0.007	1.99	1.27	1860.698	11.941	0.008	4.13	1.33
1154.734	12.103	0.006	2.32	1.27	1860.771	11.945	0.008	3.77	1.27
1154.773	12.095	0.006	2.93	1.30	1860.844	11.957	0.006	3.60	1.30
1154.840	12.083	0.005	3.03	1.42	1861.612	12.199	0.009	3.77	1.56
1155.648	11.909	0.007	2.47	1.31	1861.674	12.229	0.005	3.75	1.38
1155.705	11.900	0.010	1.87	1.27	1861.803	12.285	0.003	3.47	1.27
1155.741	11.899	0.008	1.81	1.28	1861.849	12.300	0.005	3.48	1.32
1155.770	11.893	0.007	1.83	1.30	1862.583	12.363	0.021	4.06	1.67
1155.820	11.883	0.005	2.09	1.38	1862.644	12.351	0.006	2.92	1.44
1155.846	11.876	0.003	2.57	1.45	1862.711	12.322	0.008	3.37	1.31
1499.849	11.964	0.005	2.88	1.33	1862.769	12.302	0.004	2.92	1.27
1500.607	12.104	0.009	4.09	1.53	1862.847	12.284	0.005	3.58	1.31
1500.671	12.128	0.005	3.44	1.36	1863.568	12.082	0.005	2.57	1.73
1500.742	12.155	0.006	3.26	1.27	1863.744	12.045	0.007	2.94	1.28
1500.812	12.174	0.006	4.07	1.29	1863.856	12.014	0.005	2.80	1.33
1500.853	12.199	0.006	3.14	1.34	1864.605	11.890	0.007	1.85	1.55
1501.559	12.405	0.007	2.82	1.74	1864.666	11.891	0.005	1.93	1.37
1501.633	12.397	0.006	3.16	1.44	1864.713	11.874	0.006	2.13	1.30
1501.686	12.395	0.004	3.26	1.33	1864.791	11.872	0.005	2.67	1.27
1501.777	12.406	0.005	2.89	1.27	2212.743	11.986	0.007	2.96	1.32
1501.821	12.411	0.006	2.78	1.30	2212.808	11.976	0.006	2.59	1.27
1501.856	12.400	0.005	2.82	1.35	2212.857	11.964	0.006	2.89	1.28
1502.658	12.138	0.007	2.78	1.37	2213.679	11.867	0.006	2.56	1.45
1502.741	12.120	0.005	2.00	1.27	2213.740	11.861	0.007	2.18	1.32
1502.814	12.110	0.005	2.14	1.29	2213.787	11.860	0.005	2.50	1.27
1502.848	12.098	0.006	2.28	1.34	2213.843	11.848	0.008	2.49	1.27
1503.672	11.973	0.005	2.56	1.34	2214.663	11.859	0.006	1.96	1.50
1503.735	11.971	0.006	2.54	1.27	2214.724	11.870	0.019	2.92	1.34
1503.789	11.959	0.005	2.94	1.28	2214.725	11.848	0.020	3.09	1.34
1503.811	11.961	0.008	2.90	1.29	2214.772	11.863	0.025	2.57	1.28
1503.827	11.949	0.004	3.14	1.31	2214.804	11.881	0.006	2.18	1.27
1503.846	11.946	0.006	2.88	1.34	2214.840	11.893	0.006	2.07	1.27
1504.565	11.878	0.005	3.63	1.67	2214.864	11.890	0.006	2.27	1.29

although the variation of the shape of the secondary eclipse between 1998 and 2000 increases the uncertainties of the solution.

Unfortunately, it was not possible to derive a photometric q from our light curve. We found solutions with q ($q = M_2/M_1$, where M_2 corresponds to the B supergiant) ranging from 0.6 to 1.67, with practically the same degree of significance. The solution with $q = 1.67$ gives a mass of

$\sim 127 M_\odot$ for the B0 I star, therefore can be disregarded. Only the solutions with lower mass ratio ($q \sim 0.7 \sim 0.8$) provide masses and radii according with what is expected for B supergiants. However, for all cases, the solutions correspond to overcontact systems. In these configurations, thermal contact between both components exists, hence the surface temperatures of the stars are similar. It is compatible with the light curve, that shows eclipses of similar depth, but




Figure 2. Normalised CCD spectra of Sk-69°194 obtained at CASLEO, corresponding (from top to bot) to phases 0.016, 0.284, 0.503 and 0.634, respectively.

the very different spectra of the stars suggest unlike physical conditions of their atmospheres. In spite of the wide range of mass ratios of the possible solutions, the orbital inclination is $75 \pm 1^\circ$ for all of them. This inclination, together with the observed radial velocity curve, yields for a_2 a value of $49.1 \pm 4.3 R_\odot$.

7 DISCUSSION

Massey et al. (2000) classified Sk-69°194 as B0 I+WN, and pointed out that the weakness of the WR emissions presumably is due to the continuum being dominated by the B supergiant. The temperature inferred from the light curve is compatible with that of a late WN, but all the solutions point to comparable magnitudes. In addition, no orbital movement is detected in the emission, although it would be intrinsically difficult to measure due to its weakness.

On weighing the observational data and the W-D modelling, we speculate about the alternative nature of the companion of the B supergiant:

- May be that it is truly a late WN star, with very weak emissions. The fact that no orbital movement is detectable from HeII 4686 can be ascribed to the difficulty of measuring the radial velocity of a broad emission superposed with the absorption lines of the other star.
- It is possible that we are seeing a thick disk that hide one of the binary components. The HeII 4686 emission can be originated in the region between (or surrounding) the stars.

Both hypothesis, though, do not agree with the overcontact configuration inferred from the W-D solutions. In one case, this contradiction can be ascribed to the interaction effects caused by the WR winds and radiation pressure (see, for example, Drechsel et al. 1995). On the other hand, if we are seeing an accretion disk, the W-D model must be considered with extreme caution, since the system's shape would be very different from the Roche-lobe geometry. Kondo et al. (2002) reviewed the main problems of close binary modelling, especially that concerning to Roche-lobe geometry. Sk-69°194 is a system where such objections appear to be especially pertinent. Anyhow, it is interesting to mention




Figure 3. Observed and modelled radial velocity curve for Sk-69°194 and their O-C residuals.

that Walborn et al. (2002) suggest three different ways that leads to the WN stage, one of which being mass transfer binaries. In fact, the lack of discernible spectral features of the B companion can be explained if we suppose that it is behind a thick stream of turbulent matter, hypothesis that can also give account for the eclipse shape variation occurred between 1998 and 2001. A spectroscopic study encompassing a complete orbital period, at a higher resolution than that one can reach at CASLEO, could contribute to enlighten this issue.

The system remarkably resembles HD 163181 ($\equiv$ V453 Scorpii) (Josephs et al. 2001, and references there in), in which only one spectrum is visible (BN0.5 Ia). That system has a period of ~ 12 days with radial velocity excursions of nearly 400 km s^{-1} , also exhibiting enhanced nitrogen abundance and signs of mass transfer.

From the derived values for a_2 and i , we have

$$a = 49.1(1 + q) R_{\odot}$$

$$M_1 = 10.65(1 + q)^2 M_{\odot}$$

and

$$M_2 = 10.65q(1 + q)^2 M_{\odot}.$$

Assuming for the B0 supergiant a mass of $\sim 25 M_{\odot}$ (Schmidt-Kaler 1982), the corresponding q is ~ 0.75 and M_1 is $\sim 32.5 M_{\odot}$. However, it is important to have in mind that the derived value for the inclination depends on the assumption of Roche-lobe geometry, although its constant value for a wide range of possible mass ratios suggests that it should not be very far from the actual value.

ACKNOWLEDGMENTS

I would like especially to thank Dr. Claudia Giordano for helping me with English. I also wish to acknowledge Mariela Corti for the acquisition of two of the spectra, as an interchange of telescope time. I also want to thank Dr. Nidia Morrell and Dr. Rodolfo Barbá for helpful comments, and Dr. Virpi Niemelä for her power of motivation.

The author acknowledges the use of the CCD and data acquisition system at CASLEO supported through a U.S. National Science Foundation grant AST-90-15827 to R. M. Rich. The focal reducer in use at CASLEO was kindly provided by Dr. M. Shara.

REFERENCES

- Drechsel H., Haas S., Lorenz R., Gayler S., 1995, *A&A*, 294, 723
- Isserstedt J., 1975, *A&A Suppl.*, 19, 259
- Josephs T.S., Gies D.R., Bagnuolo W.G., Shure M.A., Penny L.R., Wang Z., *PASP* 113, 957
- Kondo Y., McCluskey Jr. G.E., Guinan E.F., 2002, *New Astron. Reviews*, 46, 1
- Landolt A.U., 1992, *AJ* 104, 340
- Lucke P.B., Hodge P.W., 1970, *AJ*, 75, 171
- Massey P., Waterhouse E., De Gioia-Eastwood K., *AJ*, 119, 2214
- Sanduleak, N. 1969, *Cerro Tololo Inter-American Obs. Contr. No.* 89
- Schmidt-Kaler, Th., 1982, in: *Landolt-Börnstein, Numerical Data and Functional Relationships in Science and Technology*, New Series, ed. K. Shaifers & H.H. Voigt, Group VI, Vol. 2/b, Springer-Verlag, Berlin-Heidelberg
- Stetson P.B., 1987, *PASP* 99, 191
- Stetson P.B., 1991, In: *Grosbøl P.J., Warmel R.H. (eds.)*
- Walborn N.R., Fitzpatrick E.L., 1990, *PASP* 102, 650
- Walborn N.R., Howarth I.D., Lennon D.J., Massey P., Oey M.S., Moffat A.F.J., Skalkowski G., Morrell N.I., Drissen L., & Parker J. Wm., 2002, *AJ* 123, 2754
- Westerlund B.E., 1961, *Uppsala Astron. Obs. Ann.*, 5, 1
- Wilson R.E., 1990, *ApJ* 356, 613
- Wilson R.E., 1993, in *New Frontiers in Binary Star Research*, ed. K.C. Leung & I.S. Nha, *ASP Conf. Ser.*, 38, 91
- Wilson R.E., Devinney E.J., 1971, *ApJ* 166, 605

This paper has been typeset from a $\text{\TeX}$ / $\text{\LaTeX}$ file prepared by the author.